



Occupational Injuries and Hazards Faced by Orthopaedic Surgeons

Occupational injuries and hazards faced by orthopaedic surgeons have gained increased attention within published literature.² Occupational injuries may result in lost productivity due to surgeon absenteeism or presenteeism and may impact the quality of surgical care offered to the patient due to performance issues the surgeon may face while dealing with or recovering from the injury.^{2,3} The data below are from published literature on occupational injuries and hazards faced by orthopaedic surgeons.

Musculoskeletal Disorders

- Operating can lead surgeons to adopt postures and repetitive motions that have damaging long-term effects on physical health through imparting musculoskeletal fatigue, damaging a surgeon's muscles, nerves, and/or joints^{1,4}
- May result in chronic pain for the surgeon, which may lead to other implications including productivity loss.¹

Workplace Injuries



- Many orthopaedic surgeons sustain occupational injuries during their careers.^{2,3}
- The volume of work days missed suggests that occupational injury has economic implications for the healthcare system and providers.^{2,3}
- Occupational injury may impact the quality of surgical care offered to the patient due to performance issues the surgeon may face while dealing with or recovering from the injury.^{2,3}

Emotional and Psychological Health

- Mental health issues and other psychological problems (including anxiety, chronic fatigue, and burnout) are often dismissed as the surgeons do not wish to let their colleagues down.⁵
- There is a misconception that surgeons better cope with stress and are protected from burnout.⁵

IMPACT OF MUSCULOSKELETAL INJURIES ON ORTHOPAEDIC SURGEONS

Prevalence of the most common musculoskeletal disorders among respondent arthroplasty surgeons.⁶



orthopaedic surgeons
developed a musculoskeletal
disorder from work⁶

Productivity Loss From Occupational Injuries and Hazards for Orthopaedic Surgeons in the US.

ABSENTEEISM

Work absence due to injury or illness

22-27% OF INJURED
ORTHOPAEDIC SURGEONS

required time off from work due to a work-related injury (mean= 7.3 days [standard deviation:21.1])^{2,3,6}

4% OF INJURED
ORTHOPAEDIC SURGEONS

required longer than 3 weeks' time off due to an injury²

SURGERY WAS
REQUIRED TO TREAT **31%**

of orthopedic surgeons suffering from work-related injuries.⁶

RESPECTIVE STUDY FOUND **STRAIN INJURIES**

of upper and lower limbs were responsible for **14.8% of work-related injuries** among healthcare workers* & resulted in **3,450 lost days of work**, with the **median duration of work absence being 10.0 days**⁷

\$3,286,764

Annual net revenue generated by an orthopaedic surgeon¹⁰

Surgeon Absenteeism

(mean= 7.3 days [standard deviation:21.1])³

\$71,000-\$147,000*

Lost revenue for the institution^{2,3,10}

**DEPUY SYNTHES OFFERS TECHNOLOGY THAT MAY HELP ADDRESS OCCUPATIONAL INJURIES FACED BY
ORTHOPAEDIC SURGEONS. CONTACT YOUR LOCAL DEPUY SYNTHES SALES CONSULTANT TO LEARN MORE.**

* Healthcare workers include medical staff (eg, medical specialists or residents), registered nursing personnel, nursing assistants, healthcare technicians (eg, laboratory and radiology technicians), administrative assistants, and other professional groups (eg, hospital managers).

* Presented in 2022 USD value, inflated using Consumer Price Index

Please refer to the instructions for use for a complete list of indications, contraindications, warnings, and precautions.
The third-party trademarks used herein are the trademarks of their respective owners.



KINCISE™
Surgical Automated System

www.deploysynthes.com

© DePuy Synthes 2020, 2023. All rights reserved.
142242-221130 DSUS/EMEA

References: 1. Epstein S, Sparer EH, Tran BN, Ruan QZ, Dennerlein JT, Singh D, Lee BT. Prevalence of work-related musculoskeletal disorders among surgeons and interventionalists. *JAMA Surgery*. 2017;153(2). 2. Davis WT, Sathiyakumar V, Jahangir AA, Obremskey WT, Sethi MK. Occupational injury among orthopaedic surgeons. *J Bone Joint Surg Am*. 2013;95(16):e107. 3. Davis WT, Fletcher SA, Guillamondegui OD. Musculoskeletal occupational injury among surgeons: effects for patients, providers, and institutions. *J Surg Res*. 2014;189(2):207-212.e6. 4. Lester JD, Hsu S, Ahmad CS. Occupational hazards facing orthopaedic surgeons. *Am J Orthop*. 2012;41(3):132-139. 5. Parry DA, Oeppen RS, Amin MSA, Brennan PA. Could exercise improve mental health and cognitive skills for surgeons and other healthcare professionals? *Br J Oral Maxillofac Surg*. 2018;56(5):367-370. 6. Alqahtani SM, Alzahrani MM, Tanzer M. Adult reconstructive surgery: a high-risk profession for work-related injuries. *J Arthroplasty*. 2016;31(6):1194-1198. 7. Amaro J, Magalhaes J, Leite M, et al. Musculoskeletal injuries and absenteeism among healthcare professionals-ICD-10 characterization. *PLoS One*. 2018;13(12):e0207837. 8. Soueid A, Oudit D, Thiagarajah S, Laitung G. The pain of surgery: pain experienced by surgeons while operating. *Int J Surg*. 2010;8(2):118-120. 9. Knudsen ML, Ludewig PM, Braman JP. Musculoskeletal pain in resident orthopaedic surgeons: results of a novel survey. *Iowa Orthop J*. 2014;34:190-196. 10. Merritt Hawkins. Physician Inpatient/Outpatient Revenue Survey. 2019. www.merrithawkins.com/uploadedFiles/MerrittHawkins_RevenueSurvey_2019.pdf